

Universality in Anisotropic Linear Anelasticity*

Arash Yavari^{†1,2} and Alain Goriely³

¹*School of Civil and Environmental Engineering, Georgia Institute of Technology, Atlanta, GA 30332, USA*

²*The George W. Woodruff School of Mechanical Engineering, Georgia Institute of Technology, Atlanta, GA 30332, USA*

³*Mathematical Institute, University of Oxford, Oxford, OX2 6GG, UK.*

June 16, 2022

Abstract

In linear elasticity, universal displacements for a given symmetry class are those displacements that can be maintained by only applying boundary tractions (no body forces) and for arbitrary elastic constants in the symmetry class. In a previous work, we showed that the larger the symmetry group, the larger the space of universal displacements. Here, we generalize these ideas to the case of anelasticity. In linear anelasticity, the total strain is additively decomposed into elastic strain and anelastic strain, often referred to as an eigenstrain. We show that the *universality constraints* (equilibrium equations and arbitrariness of the elastic constants) completely specify the *universal elastic strains* for each of the eight anisotropy symmetry classes. The corresponding universal eigenstrains are the set of solutions to a system of second-order linear PDEs that ensure compatibility of the total strains. We show that for three symmetry classes, namely triclinic, monoclinic, and trigonal, only compatible (impotent) eigenstrains are universal. For the remaining five classes universal eigenstrains (up to the impotent ones) are the set of solutions to a system of linear second-order PDEs with certain arbitrary forcing terms that depend on the symmetry class.

Keywords: Universal deformation, universal displacement, linear elasticity, anelasticity, anisotropic solids, eigenstrain.

1 Introduction

In nonlinear anelasticity, in the notion first defined in [Eckart, 1948], strain has an elastic and an anelastic part. In terms of deformation gradient it is written as $\mathbf{F} = \mathbf{F}^e \mathbf{F}^a$, where \mathbf{F}^e and \mathbf{F}^a are the elastic and anelastic deformation tensors, respectively [Sadik and Yavari, 2017, Goodbrake et al., 2020a, Sozio and Yavari, 2020]. The hybrid German-English portmanteau term *eigenstrain* has its origin in the pioneering paper of Hans Reissner [Reissner, 1931] (Eigenspannung means proper or self strain) and was further popularized by Mura [Kinoshita and Mura, 1971, Mura, 1982]. In the literature several equivalent terms have been used for the same concept; *initial strain* [Kondo, 1949], *nuclei of strain* [Mindlin and Cheng, 1950], *transformation strain* [Eshelby, 1957], *inherent strain* [Ueda et al., 1975], and *residual strains* [Ambrosi et al., 2019] (see also [Jun and Korsunsky, 2010, Zhou et al., 2013]). In the setting of linear elasticity, and for infinite bodies, inclusions and their induced stress fields were systematically studied by Eshelby in a celebrated paper [Eshelby, 1957]. He showed that an ellipsoidal inclusion that has uniform eigenstrain and is embedded in an infinite linear elastic medium, has a uniform stress field. It is known that in the case of finite bodies the stress field of an inclusion with uniform eigenstrain is not necessarily uniform, e.g., a spherical inclusion

*To appear in the *Journal of Elasticity*.

[†]Corresponding author, e-mail: arash.yavari@ce.gatech.edu

centered at a finite ball [Li et al., 2007]. The extension of Eshelby’s analysis of eigenstrains to nonlinear anelasticity has received attention in the last twenty years. In the case of some special constitutive equations one can mention [Ru and Schiavone, 1996, Diani and Parks, 2000, Ru et al., 2005, Kim and Schiavone, 2007, 2008, Kim et al., 2008]. There are several more recent works that use geometric techniques [Yavari, 2013, Yavari and Goriely, 2015a,b, Golgoon et al., 2016, Golgoon and Yavari, 2017, 2018, Yavari, 2021b].

If one is asked to deform an elastic body to a desired arbitrary shape, most likely body forces will be required to achieve the desired deformation, especially when one does not specify a particular material. However, there are special deformations that can be maintained by applying only boundary tractions for any member of a material class. These are called *universal deformations*. The systematic study of universal deformations began in two seminal papers of Ericksen [Ericksen, 1954, 1955] motivated by earlier works of Rivlin [Rivlin, 1948, 1949a,b]. Ericksen proved that for compressible isotropic solids only homogeneous deformations are universal. For incompressible isotropic solids he found four families of universal deformations, in addition to the isochoric homogeneous deformations. He conjectured that only homogeneous deformations have constant principal invariants. This turned out to be incorrect [Fosdick, 1966], and led to the discovery of a fifth family of universal deformations [Singh and Pipkin, 1965, Klingbeil and Shield, 1966]. Existence of other constant principal invariant inhomogeneous universal deformations is still an open problem, but the current conjecture is that none exists.

We extended Ericksen’s analysis to compressible anelasticity and showed that universal deformations must be covariantly homogeneous [Yavari and Goriely, 2016]. We proved that this implies that for simply-connected bodies universal eigenstrains are impotent. Universal deformations and eigenstrains in incompressible anelasticity were investigated in [Goodbrake et al., 2020b]. It was shown that the six known families of universal deformations in incompressible isotropic elasticity are invariant under certain Lie subgroups of the special Euclidean group. In the analysis of universal eigenstrains it was assumed that for each class of universal deformations the corresponding universal eigenstrains have the same symmetries. Under this assumption, the universal eigenstrains were characterized for each class.

Ericksen’s analysis was extended to inhomogeneous compressible and incompressible isotropic solids in [Yavari, 2021a] (this was motivated by an earlier result in [Golgoon and Yavari, 2021]). It was shown that if the energy function is assumed to be position dependent (in the reference configuration) there are some extra universality constraints, in addition to those of the corresponding homogeneous solids. The universal inhomogeneities—the form of the position dependence of energy function consistent with the universality constraints—were fully characterized for compressible isotropic solids and for the six known universal deformations of incompressible isotropic solids.

Ericksen and Rivlin [1954] presented a limited analysis of universal deformations in anisotropic solids assuming fixed material preferred directions. We extended Ericksen’s analysis to transversely isotropic, orthotropic, and monoclinic solids in both compressible and incompressible cases [Yavari and Goriely, 2021]. We showed that for compressible transversely isotropic, orthotropic, and monoclinic solids universal deformations are homogeneous and the universal material preferred directions are uniform. For each of the six known families of universal deformations in the incompressible case (that turn out to be universal for anisotropic solids as well) we assumed that the corresponding universal material preferred directions have the same symmetries as those of the universal deformations (these symmetries are encoded in the symmetries of the right Cauchy-Green strain). Under this assumption we fully characterized the universal material preferred directions for each family and material class. Recently, we completed the universal program of nonlinear hyperelastic elasticity by characterizing the universal inhomogeneities for each of the three material classes for both compressible and incompressible cases [Yavari and Goriely, 2022a].

Since the central object of linear elasticity is displacement fields rather than deformations, *universal displacements* are the natural analogue of universal deformations in linear elasticity [Truesdell, 1966, Gurtin., 1972, Yavari et al., 2020]. These are displacements that can be maintained in the absence of body forces and by applying only boundary tractions for arbitrary elastic constants in a given symmetry class. In [Yavari et al., 2020], universal displacements of linear anisotropic elasticity were fully characterized for each of the eight symmetry classes assuming that the directions of the material anisotropy are known. Recently, we extended the analysis of universal displacements to inhomogeneous anisotropic linear elasticity [Yavari and Goriely, 2022b]. It was shown that the universality constraints of inhomogeneous linear elasticity include

those of homogeneous linear elasticity. For each of the eight symmetry classes we fully characterized the universal inhomogeneities, i.e., the form of position dependence of the elastic moduli that are consistent with the universality constraints. In the present paper, we study universality in anisotropic linear anelasticity.

This paper is organized as follows. In §2 we define universal elastic strains and eigenstrains in linear anelasticity. Universal elastic strains and eigenstrains of isotropic linear anelasticity are discussed in §3. The same problems are investigated for the other seven symmetry classes (triclinic, monoclinic, tetragonal, trigonal, orthotropic, transversely isotropic, and cubic) in §4. Conclusions are given in §5.

2 Universal elastic strains and eigenstrains in linear anelasticity

In linear anelasticity, linearized strain is additively decomposed into elastic and anelastic parts: $\epsilon = \epsilon^e + \epsilon^*$, with

$$\epsilon = \frac{1}{2} (\nabla \mathbf{u} + \nabla \mathbf{u}^T), \quad (2.1)$$

where \mathbf{u} is the displacement field and ϵ^* is the *linearized eigenstrain*, which in this paper we simply refer to as *eigenstrain*. Note that, in general, ϵ^e and ϵ^* are incompatible, i.e., $\text{curl} \circ \text{curl} \epsilon^e \neq \mathbf{0}$, and $\text{curl} \circ \text{curl} \epsilon^* \neq \mathbf{0}$, where $\text{curl} \circ \text{curl}$ is the incompatibility operator. The constitutive equations read $\sigma = \mathbf{C} \cdot \epsilon^e$, or in components

$$\sigma_{ab} = C_{abcd} (u_{c|d} - \epsilon_{cd}^*), \quad (2.2)$$

where \mathbf{C} is the elasticity tensor and summation over repeated indices is assumed. Let us consider a homogeneous linear elastic body \mathcal{B} . In the Cartesian coordinates $\{x_a\}$ the body has uniform elastic constants C_{abcd} . At $\mathbf{x} \in \mathcal{B}$, the displacement and eigenstrain fields have components $u_a(\mathbf{x})$ and $\epsilon_{ab}^*(\mathbf{x})$, respectively. In the absence of body forces, the equilibrium equations read

$$\sigma_{ab,b} = C_{abcd} u_{c,db} - C_{abcd} \epsilon_{cd,b}^* = 0. \quad (2.3)$$

It is more convenient to rewrite this in terms of elastic strains as

$$C_{abcd} \epsilon_{cd,b}^e = 0, \quad (2.4)$$

which must hold for arbitrary C_{abcd} in a given symmetry class. We refer to (2.4) as the *universality constraints*.

Definition 2.1. For a given symmetry class, a strain field that satisfies $\mathbf{C} \cdot \nabla \epsilon^e = \mathbf{0}$, or in components, $C_{abcd} \epsilon_{cd,b}^e = 0$, $a = 1, 2, 3$, for all the elasticity tensors in the symmetry class, is called a *universal elastic strain*.

The *incompatibility tensor* is defined as $\mathbf{R} = \text{curl} \circ \text{curl} \epsilon$, or in components

$$R_{ij} = R_{ji} = \varepsilon_{aki} \varepsilon_{blj} \epsilon_{ab,kl}, \quad (2.5)$$

where ε_{abc} is the permutation symbol. The six bulk compatibility equations of linear elasticity are therefore given by $R_{ij} = 0$ [Yavari, 2013]. Knowing that the total strain is compatible, one concludes that $\mathbf{R}^* = -\mathbf{R}^e$. For a given symmetry class the universality constraints (2.4) determine the set of universal elastic strains as we will see in the following sections. This leads to the definition of universal eigenstrains.

Definition 2.2. For a given symmetry class, the corresponding *universal eigenstrains* are the set of solutions to the following linear partial differential equations (PDEs)

$$\varepsilon_{aki} \varepsilon_{blj} \epsilon_{ab,kl}^* = -\varepsilon_{aki} \varepsilon_{blj} \epsilon_{ab,kl}^e, \quad (2.6)$$

where ϵ^e is a universal elastic strain fields of the symmetry class. In other words, a universal eigenstrain satisfies (2.6) for at least one universal elastic strain. Note that the above PDEs determine universal eigenstrains up to compatible (impotent) eigenstrains.

3 Isotropic linear anelasticity

For isotropic solids, in a Cartesian coordinate system $\{x_a\}$, the elasticity tensor has the representation $C_{abcd} = \lambda \delta_{ab}\delta_{cd} + \mu (\delta_{ac}\delta_{bd} + \delta_{ad}\delta_{bc})$, where λ and μ are the Lamé constants. The universality constraints (2.4) for isotropic solids are simplified to read

$$\lambda \epsilon_{cc,a}^e + 2\mu \epsilon_{ab,b}^e = 0, \quad a = 1, 2, 3. \quad (3.1)$$

Note that the above identity must hold for arbitrary elastic constants λ and μ , and hence

$$\epsilon_{cc,a}^e = \epsilon_{ab,b}^e, \quad a = 1, 2, 3. \quad (3.2)$$

Therefore, universal elastic strains have constant trace and are divergence free. A divergence-free second-order tensor can be represented by: $\epsilon_{ab}^e = \varepsilon_{acm} \varepsilon_{bdn} \phi_{cd,mn}$, where $\phi_{cd} = \phi_{dc}$ is the Beltrami potential [Beltrami, 1892, Gurtin]. In other words, any constant-trace divergence-free elastic strain is universal. Note that these universal elastic strains are incompatible as one can show that

$$R_{ij} = \phi_{kl,ijkl} + \phi_{ij,kkll} - \phi_{kj,ikkll} - \phi_{ik,jkll}. \quad (3.3)$$

Proposition 3.1. *For isotropic linear elastic solids universal elastic strains are divergence free and have constant trace. The universal eigenstrains (up to impotent eigenstrains) are the set of solutions of the following linear PDEs:*

$$\varepsilon_{aki} \varepsilon_{blj} \epsilon_{ab,kl}^* = -\phi_{kl,ijkl} - \phi_{ij,kkll} + \phi_{kj,ikkll} + \phi_{ik,jkll}, \quad (3.4)$$

where $\phi_{ij} = \phi_{ji}$ are Beltrami potentials such that $\epsilon_{cc}^e = \phi_{nn,mm} - \phi_{mn,mn}$ is constant.

4 Anisotropic linear anelasticity

In the absence of eigenstrains, Yavari et al. [2020] characterized the universal displacements of linear elasticity for all the eight anisotropy classes. In this section, we extend their work to anisotropic linear anelasticity for all the eight symmetry classes: triclinic, monoclinic, tetragonal, trigonal, orthotropic, transversely isotropic, and cubic [Cowin and Mehrabadi, 1995, Chadwick et al., 2001, Ting, 2003, Cowin and Doty, 2007]. Consider a homogeneous body made of a linear elastic solid with elasticity tensor C_{abcd} that has major $C_{abcd} = C_{cdab}$ and minor symmetries $C_{abcd} = C_{bacd}$. We use the Voigt notation with the bijection $(11, 22, 33, 23, 31, 12) \leftrightarrow (1, 2, 3, 4, 5, 6)$ to write the constitutive equations as $\sigma_\alpha = c_{\alpha\beta} \epsilon_\beta$, where Greek indices run from 1 to 6. The elasticity tensor is then represented by a symmetric 6×6 stiffness matrix as

$$\mathbf{C}(\mathbf{x}) = \begin{bmatrix} c_{11} & c_{12} & c_{13} & c_{14} & c_{15} & c_{16} \\ c_{12} & c_{22} & c_{23} & c_{24} & c_{25} & c_{26} \\ c_{13} & c_{23} & c_{33} & c_{34} & c_{35} & c_{36} \\ c_{14} & c_{24} & c_{34} & c_{44} & c_{45} & c_{46} \\ c_{15} & c_{25} & c_{35} & c_{45} & c_{55} & c_{56} \\ c_{16} & c_{26} & c_{36} & c_{46} & c_{56} & c_{66} \end{bmatrix}. \quad (4.1)$$

In this notation the equilibrium equations are written as

$$\begin{bmatrix} \frac{\partial}{\partial x_1} & 0 & 0 & 0 & \frac{\partial}{\partial x_3} & \frac{\partial}{\partial x_2} \\ 0 & \frac{\partial}{\partial x_2} & 0 & \frac{\partial}{\partial x_3} & 0 & \frac{\partial}{\partial x_1} \\ 0 & 0 & \frac{\partial}{\partial x_3} & \frac{\partial}{\partial x_2} & \frac{\partial}{\partial x_1} & 0 \end{bmatrix} \begin{bmatrix} c_{11} & c_{12} & c_{13} & c_{14} & c_{15} & c_{16} \\ c_{12} & c_{22} & c_{23} & c_{24} & c_{25} & c_{26} \\ c_{13} & c_{23} & c_{33} & c_{34} & c_{35} & c_{36} \\ c_{14} & c_{24} & c_{34} & c_{44} & c_{45} & c_{46} \\ c_{15} & c_{25} & c_{35} & c_{45} & c_{55} & c_{56} \\ c_{16} & c_{26} & c_{36} & c_{46} & c_{56} & c_{66} \end{bmatrix} \begin{bmatrix} \epsilon_1^e \\ \epsilon_2^e \\ \epsilon_3^e \\ \epsilon_4^e \\ \epsilon_5^e \\ \epsilon_6^e \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \quad (4.2)$$

where

$$\epsilon_1^e = \epsilon_{11}^e, \quad \epsilon_2^e = \epsilon_{22}^e, \quad \epsilon_3^e = \epsilon_{33}^e, \quad \epsilon_4^e = 2\epsilon_{23}^e, \quad \epsilon_5^e = 2\epsilon_{13}^e, \quad \epsilon_6^e = 2\epsilon_{12}^e. \quad (4.3)$$

Equation (4.2) and the arbitrariness of the elastic constants for a given symmetry class forces the elastic strains to satisfy a set of PDEs that we call *universality constraints*.

4.1 Triclinic solids

Triclinic solids are the least symmetric among the eight symmetry classes; the identity and minus identity are the only symmetry transformations for such materials. This means that triclinic linear elastic solids have 21 independent elastic constants. In [Yavari et al., 2020] it was shown that for triclinic linear elastic solids homogeneous displacements are the only universal displacements. It is straightforward to show that for triclinic solids, the universality constraints (2.4) read $\epsilon_{\beta,a} = 0$, for $\beta = 1, \dots, 6$, and $a = 1, 2, 3$.¹ This means that universal elastic strains are constant. This implies that universal elastic strains are compatible, and consequently the universal eigenstrains are impotent.

Proposition 4.1. *For triclinic linear elastic solids, universal elastic strains are uniform, and consequently, universal eigenstrains are impotent.*

Remark 4.2. The universal eigenstrains are of the form $\epsilon^*(\mathbf{x}) = \frac{1}{2} [\nabla \mathbf{u}^*(\mathbf{x}) + \nabla \mathbf{u}^*(\mathbf{x})^\top]$, where $\mathbf{u}^*(\mathbf{x})$ is any displacement field. For a given impotent eigenstrain $\epsilon^*(\mathbf{x})$, universal displacements are superposition of $\mathbf{u}^*(\mathbf{x})$ and all homogeneous displacements. This implies that all displacement fields are universal. However, if one defines universal eigenstrains modulo impotent eigenstrains, in the case of triclinic solids, only homogeneous displacements are universal.

4.2 Monoclinic solids

In a monoclinic solid there is one plane of material symmetry that we assume to be parallel to the x_1x_2 -plane. A monoclinic linear elastic solid has 13 independent elastic constants. The elasticity matrix has the following

¹All the symbolic computations in this paper were performed using Mathematica Version 13.0.0.0, Wolfram Research, Champaign, IL.

form:

$$\mathbf{c} = \begin{bmatrix} c_{11} & c_{12} & c_{13} & 0 & 0 & c_{16} \\ c_{12} & c_{22} & c_{23} & 0 & 0 & c_{26} \\ c_{13} & c_{23} & c_{33} & 0 & 0 & c_{36} \\ 0 & 0 & 0 & c_{44} & c_{45} & 0 \\ 0 & 0 & 0 & c_{45} & c_{55} & 0 \\ c_{16} & c_{26} & c_{36} & 0 & 0 & c_{66} \end{bmatrix}. \quad (4.4)$$

In [Yavari et al., 2020] it was shown that for a monoclinic linear elastic solid with planes of symmetry parallel to the x_1x_2 -plane, universal displacements are the superposition of homogeneous displacements $\mathbf{F} \cdot \mathbf{x}$ (\mathbf{F} is a constant matrix) and the one-parameter inhomogeneous displacement field $(cx_2x_3, -cx_1x_3, 0)$.

For monoclinic solids, the universality constraints (4.2) read $\epsilon_{\beta,a} = 0$, for $\beta = 1, 2, 3, 6$, and $a = 1, 2, 3$, and

$$\begin{aligned} \frac{\partial \epsilon_{23}^e}{\partial x_2} &= \frac{\partial \epsilon_{23}^e}{\partial x_3} = 0, \\ \frac{\partial \epsilon_{13}^e}{\partial x_2} &= \frac{\partial \epsilon_{13}^e}{\partial x_3} = 0, \\ \frac{\partial \epsilon_{13}^e}{\partial x_2} + \frac{\partial \epsilon_{23}^e}{\partial x_1} &= 0. \end{aligned} \quad (4.5)$$

The first two PDEs imply that $\epsilon_{23}^e = \epsilon_{23}^e(x_1)$, and $\epsilon_{13}^e = \epsilon_{13}^e(x_2)$, and the third PDE implies that $\epsilon_{23}^e(x_1) = -\epsilon_{13}^e(x_2) = c_0$, a constant. It is straightforward to check that the universal elastic strains are compatible.

Proposition 4.3. *For monoclinic linear elastic solids the universal elastic strains have the following form*

$$\epsilon^e(\mathbf{x}) = \begin{bmatrix} c_1 & c_6 & -c_0x_2 + c_5 \\ c_6 & c_2 & c_0x_1 + c_4 \\ -c_0x_2 + c_5 & c_0x_1 + c_4 & c_3 \end{bmatrix}, \quad (4.6)$$

where $c_i, i = 0, 1, \dots, 6$ are constants. These strains are compatible, and consequently, universal eigenstrains are impotent.

Remark 4.4. The universal eigenstrains are of the form $\epsilon^*(\mathbf{x}) = \frac{1}{2} [\nabla \mathbf{u}^*(\mathbf{x}) + \nabla \mathbf{u}^*(\mathbf{x})^\top]$, where $\mathbf{u}^*(\mathbf{x})$ is any displacement field. For a given impotent eigenstrain $\epsilon^*(\mathbf{x})$, universal displacements are superposition of $\mathbf{u}^*(\mathbf{x})$ and all universal displacements in the absence of eigenstrains, i.e., those of linear elasticity [Yavari et al., 2020, Proposition 3.2]. This implies that all displacement fields are universal. However, if one defines universal eigenstrains modulo impotent eigenstrains, in the case of monoclinic solids, universal displacements are identical to those given in [Yavari et al., 2020, Proposition 3.2].

4.3 Tetragonal solids

A tetragonal solid has five symmetry planes. The normals of four of them are coplanar and the fifth plane is normal to the other four. In a Cartesian coordinate system (x_1, x_2, x_3) we assume, without loss of generality, that the fifth normal is parallel to the x_3 axis. Two of the symmetry planes are parallel to the x_1x_3 and x_2x_3 -planes. The other two symmetry planes are related to the ones parallel to the x_1x_3 -plane by $\pi/4$ and

$3\pi/4$ rotations about the x_3 axis. A tetragonal solid has 6 independent elastic constants and the elasticity matrix has the following form:

$$\mathbf{c} = \begin{bmatrix} c_{11} & c_{12} & c_{13} & 0 & 0 & 0 \\ c_{12} & c_{11} & c_{13} & 0 & 0 & 0 \\ c_{13} & c_{13} & c_{33} & 0 & 0 & 0 \\ 0 & 0 & 0 & c_{44} & 0 & 0 \\ 0 & 0 & 0 & 0 & c_{44} & 0 \\ 0 & 0 & 0 & 0 & 0 & c_{66} \end{bmatrix}. \quad (4.7)$$

In [Yavari et al., 2020] it was shown that in a tetragonal linear elastic solid with the tetragonal axes parallel to the x_3 -axis, the universal displacements are a superposition of homogeneous displacements and the following inhomogeneous displacements:²

$$\begin{aligned} u_1(x_1, x_2, x_3) &= F_{11}x_1 + F_{12}x_2 + F_{13}x_3 + c_1x_2x_3 + c_2x_1x_3, \\ u_2(x_1, x_2, x_3) &= F_{21}x_1 + F_{22}x_2 + F_{23}x_3 - c_2x_2x_3 + c_3x_1x_3, \\ u_3(x_1, x_2, x_3) &= F_{31}x_1 + F_{32}x_2 + F_{33}x_3 + g(x_1, x_2), \end{aligned} \quad (4.8)$$

where c_1 and c_2 are constants, and $g = g(x_1, x_2)$ is a harmonic function.

The universality constraints (4.2) give us the following thirteen PDEs:

$$\begin{aligned} \frac{\partial \epsilon_{11}^e}{\partial x_1} &= \frac{\partial \epsilon_{11}^e}{\partial x_2} = 0, \\ \frac{\partial \epsilon_{22}^e}{\partial x_1} &= \frac{\partial \epsilon_{22}^e}{\partial x_2} = 0, \\ \frac{\partial \epsilon_{11}^e}{\partial x_3} + \frac{\partial \epsilon_{22}^e}{\partial x_3} &= 0, \\ \frac{\partial \epsilon_{33}^e}{\partial x_1} &= \frac{\partial \epsilon_{33}^e}{\partial x_2} = \frac{\partial \epsilon_{33}^e}{\partial x_3} = 0, \\ \frac{\partial \epsilon_{12}^e}{\partial x_1} &= \frac{\partial \epsilon_{12}^e}{\partial x_2} = 0, \\ \frac{\partial \epsilon_{13}^e}{\partial x_3} &= \frac{\partial \epsilon_{23}^e}{\partial x_3} = 0, \\ \frac{\partial \epsilon_{13}^e}{\partial x_1} + \frac{\partial \epsilon_{23}^e}{\partial x_2} &= 0. \end{aligned} \quad (4.9)$$

The first five PDEs imply that $\epsilon_{11}^e = \epsilon_{11}^e(x_3)$, $\epsilon_{22}^e = -\epsilon_{11}^e(x_3) + c_0$, where c_0 is a constant. Eq.(4.9)₄ implies that ϵ_{33}^e is constant. Eq.(4.9)₅ implies that $\epsilon_{12}^e = \epsilon_{12}^e(x_3)$. The last three PDEs imply that

$$\epsilon_{13}^e = \epsilon_{13}^e(x_1, x_2), \quad \epsilon_{23}^e = \epsilon_{23}^e(x_1, x_2), \quad \text{with the constraint} \quad \frac{\partial \epsilon_{13}^e}{\partial x_1} + \frac{\partial \epsilon_{23}^e}{\partial x_2} = 0. \quad (4.10)$$

Therefore, we have the following result.

Proposition 4.5. *For a tetragonal linear elastic solid with the tetragonal axis parallel to the x_3 -axis in a*

²There is a typo in Eq.(3.22)₂ in Yavari et al. [2020]: $-c_2x_1x_3$ should read $-c_2x_2x_3$.

Cartesian coordinate system (x_1, x_2, x_3) , the universal elastic strains have the following form

$$\boldsymbol{\epsilon}^e(\mathbf{x}) = \begin{bmatrix} \epsilon_{11}^e(x_3) & \epsilon_{12}^e(x_3) & \epsilon_{13}^e(x_1, x_2) \\ \epsilon_{12}^e(x_3) & -\epsilon_{11}^e(x_3) + c_0 & \epsilon_{23}^e(x_1, x_2) \\ \epsilon_{13}^e(x_1, x_2) & \epsilon_{23}^e(x_1, x_2) & \epsilon_{33}^e \end{bmatrix}, \quad (4.11)$$

where c_0 and ϵ_{33}^e are constants, $\epsilon_{11}^e(x_3)$, $\epsilon_{12}^e(x_3)$, and $\epsilon_{13}^e(x_1, x_2)$ are arbitrary functions, and $\epsilon_{23}^e(x_1, x_2)$ has the following representation

$$\epsilon_{23}^e(x_1, x_2) = - \int \epsilon_{13,1}^e(x_1, x_2) dx_2 + f(x_1), \quad (4.12)$$

for an arbitrary function $f(x_1)$. Universal eigenstrains (up to impotent eigenstrains) are solutions of the six second-order linear PDEs $\text{curl} \circ \text{curl} \boldsymbol{\epsilon}^* = -\mathbf{R}^e$.

Remark 4.6. The incompatibility tensor of the universal elastic strains reads

$$\mathbf{R}^e = \begin{bmatrix} -\epsilon_{11}^{e''}(x_3) & \epsilon_{12}^{e''}(x_3) & -\nabla^2 \epsilon_{13}^e(x_1, x_2) \\ \epsilon_{12}^{e''}(x_3) & \epsilon_{11}^{e''}(x_3) & -\nabla^2 \epsilon_{23}^e(x_1, x_2) \\ -\nabla^2 \epsilon_{13}^e(x_1, x_2) & -\nabla^2 \epsilon_{23}^e(x_1, x_2) & 0 \end{bmatrix}. \quad (4.13)$$

It is seen that the universal elastic strains (and consequently the universal eigenstrains) are impotent if ϵ_{11}^e and ϵ_{12}^e are linear functions, and ϵ_{13}^e and ϵ_{23}^e are harmonic.

4.4 Trigonal solids

A trigonal solid has three planes of symmetry whose normals lie in the same plane and are related by $\pi/3$ rotations. We choose Cartesian coordinates (x_1, x_2, x_3) such that the trigonal axis is parallel to the x_3 -axis. A trigonal solid has 6 independent elastic constants and the elasticity matrix has the following form:

$$\mathbf{c} = \begin{bmatrix} c_{11} & c_{12} & c_{13} & 0 & c_{15} & 0 \\ c_{12} & c_{11} & c_{13} & 0 & -c_{15} & 0 \\ c_{13} & c_{13} & c_{33} & 0 & 0 & 0 \\ 0 & 0 & 0 & c_{44} & 0 & -c_{15} \\ c_{15} & -c_{15} & 0 & 0 & c_{44} & 0 \\ 0 & 0 & 0 & -c_{15} & 0 & \frac{1}{2}(c_{11} - c_{12}) \end{bmatrix}. \quad (4.14)$$

Yavari et al. [2020] showed that in the absence of eigenstrains universal displacements are a superposition of homogeneous displacements and the following inhomogeneous displacements

$$\begin{aligned} u_1^{\text{inh}}(x_1, x_2, x_3) &= a_{123}x_1x_2x_3 + a_{12}x_1x_2 + a_{13}x_1x_3 + a_{23}x_2x_3, \\ u_2^{\text{inh}}(x_1, x_2, x_3) &= \frac{1}{2}(a_{12} + a_{123}x_3)(x_1^2 - x_2^2) + b_{13}x_1x_3 - a_{13}x_2x_3, \\ u_3^{\text{inh}}(x_1, x_2, x_3) &= -a_{123}x_1^2x_2 - (a_{23} + b_{13})x_1x_2 + \frac{1}{3}a_{123}x_2^3 - a_{13}(x_1^2 - x_2^2). \end{aligned} \quad (4.15)$$

The universality constraints (4.2) give us the following fourteen PDEs:

$$\begin{aligned}\frac{\partial \epsilon_{33}^e}{\partial x_1} &= \frac{\partial \epsilon_{33}^e}{\partial x_2} = \frac{\partial \epsilon_{33}^e}{\partial x_3} = 0, \\ \frac{\partial \epsilon_{13}^e}{\partial x_3} &= \frac{\partial \epsilon_{23}^e}{\partial x_3} = 0,\end{aligned}\tag{4.16}$$

and

$$\begin{aligned}\frac{\partial \epsilon_{11}^e}{\partial x_3} + \frac{\partial \epsilon_{22}^e}{\partial x_3} &= 0, \\ \frac{\partial \epsilon_{11}^e}{\partial x_1} + \frac{\partial \epsilon_{12}^e}{\partial x_2} &= 0, \\ \frac{\partial \epsilon_{22}^e}{\partial x_1} - \frac{\partial \epsilon_{12}^e}{\partial x_2} &= 0, \\ \frac{\partial \epsilon_{12}^e}{\partial x_1} + \frac{\partial \epsilon_{22}^e}{\partial x_2} &= 0, \\ \frac{\partial \epsilon_{12}^e}{\partial x_1} - \frac{\partial \epsilon_{11}^e}{\partial x_2} &= 0,\end{aligned}\tag{4.17}$$

and

$$\begin{aligned}\frac{\partial \epsilon_{13}^e}{\partial x_1} + \frac{\partial \epsilon_{23}^e}{\partial x_2} &= 0, \\ \frac{\partial \epsilon_{11}^e}{\partial x_3} - \frac{\partial \epsilon_{22}^e}{\partial x_3} - 2\frac{\partial \epsilon_{23}^e}{\partial x_2} + 2\frac{\partial \epsilon_{13}^e}{\partial x_2} &= 0, \\ \frac{\partial \epsilon_{11}^e}{\partial x_1} - \frac{\partial \epsilon_{22}^e}{\partial x_1} - 2\frac{\partial \epsilon_{12}^e}{\partial x_2} &= 0, \\ \frac{\partial \epsilon_{23}^e}{\partial x_1} + \frac{\partial \epsilon_{13}^e}{\partial x_2} + \frac{\partial \epsilon_{12}^e}{\partial x_3} &= 0.\end{aligned}\tag{4.18}$$

From (4.16) one concludes that ϵ_{33}^e is constant and $\epsilon_{13}^e = \epsilon_{13}^e(x_1, x_2)$, $\epsilon_{23}^e = \epsilon_{23}^e(x_1, x_2)$. From the last four PDEs in (4.17) one gets

$$\begin{aligned}\frac{\partial \epsilon_{11}^e}{\partial x_1} + \frac{\partial \epsilon_{22}^e}{\partial x_1} &= 0, \\ \frac{\partial \epsilon_{11}^e}{\partial x_2} + \frac{\partial \epsilon_{22}^e}{\partial x_2} &= 0.\end{aligned}\tag{4.19}$$

These together with (4.17)₁ imply that $\epsilon_{22}^e(x_1, x_2, x_3) = -\epsilon_{11}^e(x_1, x_2, x_3) + c_0$, where c_0 is a constant. Substituting this into (4.18)₃ one obtains

$$\frac{\partial \epsilon_{11}^e}{\partial x_1} - \frac{\partial \epsilon_{12}^e}{\partial x_2} = 0.\tag{4.20}$$

This together with (4.17)₂ implies that

$$\frac{\partial \epsilon_{11}^e}{\partial x_1} = \frac{\partial \epsilon_{12}^e}{\partial x_2} = 0,\tag{4.21}$$

and hence $\epsilon_{11}^e = \epsilon_{11}^e(x_2, x_3)$ and $\epsilon_{12}^e = \epsilon_{12}^e(x_1, x_3)$. Now the remaining PDEs are simplified to read:

$$\begin{aligned}\frac{\partial \epsilon_{13}^e}{\partial x_1} + \frac{\partial \epsilon_{23}^e}{\partial x_2} &= 0, \\ \frac{\partial \epsilon_{11}^e}{\partial x_3} - \frac{\partial \epsilon_{23}^e}{\partial x_2} + \frac{\partial \epsilon_{13}^e}{\partial x_1} &= 0, \\ \frac{\partial \epsilon_{12}^e}{\partial x_1} - \frac{\partial \epsilon_{11}^e}{\partial x_2} &= 0, \\ \frac{\partial \epsilon_{23}^e}{\partial x_1} + \frac{\partial \epsilon_{13}^e}{\partial x_2} + \frac{\partial \epsilon_{12}^e}{\partial x_3} &= 0.\end{aligned}\tag{4.22}$$

From (4.22)₃, one obtains

$$\frac{\partial}{\partial x_1} \epsilon_{12}^e(x_1, x_3) = \frac{\partial}{\partial x_2} \epsilon_{11}^e(x_2, x_3) \Rightarrow \frac{\partial^2}{\partial x_1^2} \epsilon_{12}^e(x_1, x_3) = \frac{\partial^2}{\partial x_2^2} \epsilon_{11}^e(x_2, x_3) = 0. \quad (4.23)$$

Thus, one has

$$\begin{aligned} \epsilon_{11}^e(x_2, x_3) &= x_2 f_{11}(x_3) + g_{11}(x_3), \\ \epsilon_{12}^e(x_1, x_3) &= x_1 f_{12}(x_3) + g_{12}(x_3). \end{aligned} \quad (4.24)$$

From (4.22)₃, one concludes that $f_{12}(x_3) = f_{11}(x_3)$. Equation (4.22)₂ can be rewritten as

$$\frac{\partial \epsilon_{13}^e(x_1, x_2)}{\partial x_1} - \frac{\partial \epsilon_{23}^e(x_1, x_2)}{\partial x_2} = -x_2 f'_{11}(x_3) - g'_{11}(x_3). \quad (4.25)$$

Therefore, $x_2 f''_{11}(x_3) + g''_{11}(x_3) = 0$, which implies that $f''_{11}(x_3) = g''_{11}(x_3) = 0$. Thus, $f_{11}(x_3) = a_{11}x_3 + a_0$, and $g_{11}(x_3) = b_{11}x_3 + c_1$. Similarly, from (4.22)₄ one concludes that $g_{12}(x_3) = b_{12}x_3 + c_2$.

Finally, the three PDEs in (4.22) are simplified to read

$$\begin{aligned} \frac{\partial \epsilon_{13}^e}{\partial x_1} + \frac{\partial \epsilon_{23}^e}{\partial x_2} &= 0, \\ \frac{\partial \epsilon_{13}^e}{\partial x_1} - \frac{\partial \epsilon_{23}^e}{\partial x_2} &= -a_{11}x_2 - b_{11}, \\ \frac{\partial \epsilon_{23}^e}{\partial x_1} + \frac{\partial \epsilon_{13}^e}{\partial x_2} &= -a_{11}x_1 - b_{12}. \end{aligned} \quad (4.26)$$

From the first two one obtains

$$\begin{aligned} \epsilon_{13}^e(x_1, x_2) &= -\frac{1}{2}a_{11}x_1x_2 - \frac{1}{2}b_{11}x_1 + f_{13}(x_2), \\ \epsilon_{23}^e(x_1, x_2) &= \frac{1}{4}a_{11}x_2^2 + \frac{1}{2}b_{11}x_2 + f_{23}(x_1). \end{aligned} \quad (4.27)$$

Substituting these back into (4.26)₃, one obtains $f'_{23}(x_1) + \frac{1}{2}a_{11}x_1 + b_{12} = -f'_{13}(x_2)$. This implies that $f'_{23}(x_1) + \frac{1}{2}a_{11}x_1 + b_{12} = -f'_{13}(x_2) = c_3$, and hence $f_{23}(x_1) = -\frac{1}{4}a_{11}x_1^2 + (-b_{12} + c_3)x_1 + c_4$, and $f_{13}(x_2) = -c_3x_2 + c_5$.

In summary, we have proved the following result.

Proposition 4.7. *For trigonal linear elastic solids whose trigonal axes are parallel to the x_3 axis in a Cartesian coordinate system (x_1, x_2, x_3) , the universal elastic strains have the following components*

$$\begin{aligned} \epsilon_{11}^e(x_1, x_2, x_3) &= a_{11}x_2x_3 + a_0x_2 + b_{11}x_3 + c_1, & \epsilon_{22}^e(x_1, x_2, x_3) &= -\epsilon_{11}^e(x_1, x_2, x_3) + c_0, \\ \epsilon_{33}^e(x_1, x_2, x_3) &= \epsilon_{23}^e, & \epsilon_{23}^e(x_1, x_2, x_3) &= \frac{1}{4}a_{11}x_2^2 + \frac{1}{2}b_{11}x_2 - \frac{1}{4}a_{11}x_1^2 + (-b_{12} + c_3)x_1 + c_4, \\ \epsilon_{13}^e(x_1, x_2, x_3) &= -\frac{1}{2}a_{11}x_1x_2 - \frac{1}{2}b_{11}x_1 - c_3x_2 + c_5, & \epsilon_{12}^e(x_1, x_2, x_3) &= a_{11}x_1x_3 + b_{12}x_3 + c_2. \end{aligned} \quad (4.28)$$

These strains are compatible, and consequently, universal eigenstrains are impotent.

Remark 4.8. The universal eigenstrains are of the form $\epsilon^*(\mathbf{x}) = \frac{1}{2} [\nabla \mathbf{u}^*(\mathbf{x}) + \nabla \mathbf{u}^{*\top}(\mathbf{x})]$, where $\mathbf{u}^*(\mathbf{x})$ is any displacement field. For a given impotent eigenstrain $\epsilon^*(\mathbf{x})$, universal displacements are superposition of $\mathbf{u}^*(\mathbf{x})$ and all universal displacements in the absence of eigenstrains, i.e., those of linear elasticity [Yavari et al., 2020, Proposition 3.4]. This implies that all displacement fields are universal. However, if one defines universal eigenstrains modulo impotent eigenstrains, in the case of trigonal solids, universal displacements are identical to those given in [Yavari et al., 2020, Proposition 3.4].

4.5 Orthotropic solids

An orthotropic solid has three mutually orthogonal symmetry planes. Let us choose Cartesian coordinates (x_1, x_2, x_3) whose coordinate planes are parallel to the symmetry planes. An orthotropic solid has 9 independent elastic constants, and the elasticity matrix has the following form:

$$\mathbf{c} = \begin{bmatrix} c_{11} & c_{12} & c_{13} & 0 & 0 & 0 \\ c_{12} & c_{22} & c_{23} & 0 & 0 & 0 \\ c_{13} & c_{23} & c_{33} & 0 & 0 & 0 \\ 0 & 0 & 0 & c_{44} & 0 & 0 \\ 0 & 0 & 0 & 0 & c_{55} & 0 \\ 0 & 0 & 0 & 0 & 0 & c_{66} \end{bmatrix}. \quad (4.29)$$

In [Yavari et al., 2020] it was shown that in an orthotropic linear elastic solid whose planes of symmetry are normal to the coordinate axes in a Cartesian coordinate system (x_1, x_2, x_3) , and in the absence of eigenstrains the universal displacements are the superposition of homogeneous displacement fields and the 3-parameter inhomogeneous displacement field $(a_1 x_2 x_3, a_2 x_1 x_3, a_3 x_1 x_2)$.

The universality constraints (4.2) give us the following fifteen PDEs:

$$\begin{aligned} \frac{\partial \epsilon_{11}^e}{\partial x_1} &= \frac{\partial \epsilon_{11}^e}{\partial x_2} = \frac{\partial \epsilon_{11}^e}{\partial x_3} = 0, \\ \frac{\partial \epsilon_{22}^e}{\partial x_1} &= \frac{\partial \epsilon_{22}^e}{\partial x_2} = \frac{\partial \epsilon_{22}^e}{\partial x_3} = 0, \\ \frac{\partial \epsilon_{33}^e}{\partial x_1} &= \frac{\partial \epsilon_{33}^e}{\partial x_2} = \frac{\partial \epsilon_{33}^e}{\partial x_3} = 0, \\ \frac{\partial \epsilon_{12}^e}{\partial x_1} &= \frac{\partial \epsilon_{12}^e}{\partial x_2} = 0, \\ \frac{\partial \epsilon_{13}^e}{\partial x_1} &= \frac{\partial \epsilon_{13}^e}{\partial x_3} = 0, \\ \frac{\partial \epsilon_{23}^e}{\partial x_2} + \frac{\partial \epsilon_{23}^e}{\partial x_3} &= 0. \end{aligned} \quad (4.30)$$

Thus, the normal elastic strains are constant and $\epsilon_{23}^e = \epsilon_{23}^e(x_1)$, $\epsilon_{13}^e = \epsilon_{13}^e(x_2)$, and $\epsilon_{12}^e = \epsilon_{12}^e(x_3)$. Therefore, we have proved the following result.

Proposition 4.9. *For orthotropic linear elastic solids with planes of symmetry normal to the coordinate axes in a Cartesian coordinate system (x_1, x_2, x_3) , the universal elastic strains have the following form*

$$\boldsymbol{\epsilon}^e(\mathbf{x}) = \begin{bmatrix} \epsilon_{11}^e & \epsilon_{12}^e(x_3) & \epsilon_{13}^e(x_2) \\ \epsilon_{12}^e(x_3) & \epsilon_{22}^e & \epsilon_{23}^e(x_1) \\ \epsilon_{13}^e(x_2) & \epsilon_{23}^e(x_1) & \epsilon_{33}^e \end{bmatrix}, \quad (4.31)$$

where $\epsilon_{11}^e, \epsilon_{22}^e, \epsilon_{33}^e$ are constant, and $\epsilon_{23}^e(x_1), \epsilon_{13}^e(x_2), \epsilon_{12}^e(x_3)$ are arbitrary functions. Universal eigenstrains

(up to impotent eigenstrains) are the set of solutions to the following six PDEs

$$\begin{aligned}
& \frac{\partial^2 \epsilon_{11}^*}{\partial x_2^2} + \frac{\partial^2 \epsilon_{22}^*}{\partial x_1^2} - 2 \frac{\partial^2 \epsilon_{12}^*}{\partial x_1 \partial x_2} = 0, \\
& \frac{\partial^2 \epsilon_{11}^*}{\partial x_3^2} + \frac{\partial^2 \epsilon_{33}^*}{\partial x_1^2} - 2 \frac{\partial^2 \epsilon_{13}^*}{\partial x_1 \partial x_3} = 0, \\
& \frac{\partial^2 \epsilon_{22}^*}{\partial x_3^2} + \frac{\partial^2 \epsilon_{33}^*}{\partial x_2^2} - 2 \frac{\partial^2 \epsilon_{23}^*}{\partial x_2 \partial x_3} = 0, \\
& -\frac{\partial^2 \epsilon_{11}^*}{\partial x_2 \partial x_3} - \frac{\partial^2 \epsilon_{23}^*}{\partial x_1^2} + \frac{\partial^2 \epsilon_{13}^*}{\partial x_1 \partial x_2} + \frac{\partial^2 \epsilon_{12}^*}{\partial x_1 \partial x_3} = \epsilon_{23}^e{}''(x_1), \\
& -\frac{\partial^2 \epsilon_{22}^*}{\partial x_1 \partial x_3} + \frac{\partial^2 \epsilon_{23}^*}{\partial x_1 \partial x_2} - \frac{\partial^2 \epsilon_{13}^*}{\partial x_2^2} + \frac{\partial^2 \epsilon_{12}^*}{\partial x_2 \partial x_3} = \epsilon_{13}^e{}''(x_2), \\
& -\frac{\partial^2 \epsilon_{33}^*}{\partial x_1 \partial x_2} + \frac{\partial^2 \epsilon_{23}^*}{\partial x_1 \partial x_3} + \frac{\partial^2 \epsilon_{13}^*}{\partial x_2 \partial x_3} - \frac{\partial^2 \epsilon_{12}^*}{\partial x_3^2} = \epsilon_{12}^e{}''(x_3).
\end{aligned} \tag{4.32}$$

Remark 4.10. The incompatibility tensor of the universal elastic strains reads

$$\mathbf{R}^e = \begin{bmatrix} 0 & -\epsilon_{12}^e{}''(x_3) & -\epsilon_{13}^e{}''(x_2) \\ -\epsilon_{12}^e{}''(x_3) & 0 & -\epsilon_{23}^e{}''(x_1) \\ -\epsilon_{13}^e{}''(x_2) & -\epsilon_{23}^e{}''(x_1) & 0 \end{bmatrix}. \tag{4.33}$$

We observe that the universal elastic strains (and consequently the universal eigenstrains) are impotent if $\epsilon_{23}^e(x_1), \epsilon_{13}^e(x_2), \epsilon_{12}^e(x_3)$ are linear functions.

4.6 Transversely isotropic solids

A transversely isotropic solid has an axis of symmetry that is normal to the isotropy planes. Let us assume that the axis of transverse isotropy is along the x_3 -axis in a Cartesian coordinates (x_1, x_2, x_3) . A transversely isotropic solid has 5 independent elastic constants, and the elasticity matrix has the following representation:

$$\mathbf{c} = \begin{bmatrix} c_{11} & c_{12} & c_{13} & 0 & 0 & 0 \\ c_{12} & c_{11} & c_{13} & 0 & 0 & 0 \\ c_{13} & c_{13} & c_{33} & 0 & 0 & 0 \\ 0 & 0 & 0 & c_{44} & 0 & 0 \\ 0 & 0 & 0 & 0 & c_{44} & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{1}{2}(c_{11} - c_{12}) \end{bmatrix}. \tag{4.34}$$

In [Yavari et al., 2020] it was shown that, in the absence of egenstrains, universal displacements have the following form:

$$\begin{aligned}
u_1(x_1, x_2, x_3) &= c_1 x_1 + c_2 x_2 + c x_2 x_3 + x_3 h_1(x_1, x_2) + k_1(x_1, x_2), \\
u_2(x_1, x_2, x_3) &= -c_2 x_1 + c_1 x_2 - c x_1 x_3 + x_3 h_2(x_1, x_2) + k_2(x_1, x_2), \\
u_3(x_1, x_2, x_3) &= c_3 x_3 + \hat{u}_3(x_1, x_2),
\end{aligned} \tag{4.35}$$

where $\xi(x_2 + ix_1) = h_2(x_1, x_2) + ih_1(x_1, x_2)$ and $\eta(x_2 + ix_1) = k_2(x_1, x_2) + ik_1(x_1, x_2)$ ³ are holomorphic, and $\hat{u}_3(x_1, x_2)$ is harmonic.

The universality constraints (4.2) give us the following fifteen PDEs:

$$\begin{aligned}\frac{\partial \epsilon_{33}^e}{\partial x_1} &= \frac{\partial \epsilon_{33}^e}{\partial x_2} = \frac{\partial \epsilon_{33}^e}{\partial x_3} = 0, \\ \frac{\partial \epsilon_{13}^e}{\partial x_3} &= \frac{\partial \epsilon_{23}^e}{\partial x_3} = 0, \\ \frac{\partial \epsilon_{13}^e}{\partial x_1} + \frac{\partial \epsilon_{23}^e}{\partial x_2} &= 0,\end{aligned}\tag{4.36}$$

and

$$\begin{aligned}\frac{\partial \epsilon_{11}^e}{\partial x_3} + \frac{\partial \epsilon_{22}^e}{\partial x_3} &= 0, \\ \frac{\partial \epsilon_{11}^e}{\partial x_1} + \frac{\partial \epsilon_{12}^e}{\partial x_2} &= 0, \\ \frac{\partial \epsilon_{22}^e}{\partial x_1} - \frac{\partial \epsilon_{12}^e}{\partial x_2} &= 0, \\ \frac{\partial \epsilon_{12}^e}{\partial x_1} - \frac{\partial \epsilon_{11}^e}{\partial x_2} &= 0, \\ \frac{\partial \epsilon_{12}^e}{\partial x_1} + \frac{\partial \epsilon_{22}^e}{\partial x_2} &= 0.\end{aligned}\tag{4.37}$$

From (4.36)₁ one concludes that ϵ_{33}^e is constant. The remaining PDEs in (4.36) imply that $\epsilon_{13}^e = \epsilon_{13}^e(x_1, x_2)$, $\epsilon_{23}^e = \epsilon_{23}^e(x_1, x_2)$, and $\frac{\partial \epsilon_{13}^e(x_1, x_2)}{\partial x_1} + \frac{\partial \epsilon_{23}^e(x_1, x_2)}{\partial x_2} = 0$. Therefore, $\epsilon_{13}^e(x_1, x_2)$ is an arbitrary function and

$$\epsilon_{23}^e(x_1, x_2) = - \int \epsilon_{13,1}^e(x_1, x_2) dx_2 + \hat{\epsilon}_{23}(x_1),\tag{4.38}$$

where $\hat{\epsilon}_{23}(x_1)$ is an arbitrary function. From the last four PDEs in (4.37), one obtains

$$\frac{\partial \epsilon_{11}^e}{\partial x_1} + \frac{\partial \epsilon_{22}^e}{\partial x_1} = 0, \quad \frac{\partial \epsilon_{11}^e}{\partial x_2} + \frac{\partial \epsilon_{22}^e}{\partial x_2} = 0,\tag{4.39}$$

which together with (4.37)₁ imply that $\epsilon_{22}^e(x_1, x_2, x_3) = -\epsilon_{11}^e(x_1, x_2, x_3) + c_0$, where c_0 is a constant. The remaining PDEs are (4.37)₂ and (4.37)₄. They imply that

$$\begin{aligned}\frac{\partial^2 \epsilon_{11}^e}{\partial x_1^2} + \frac{\partial^2 \epsilon_{11}^e}{\partial x_2^2} &= 0, \\ \epsilon_{12}^e(x_1, x_2, x_3) &= \int \epsilon_{11,2}^e(x_1, x_2, x_3) dx_1 + \hat{\epsilon}(x_2, x_3),\end{aligned}\tag{4.40}$$

where $\hat{\epsilon}(x_2, x_3)$ is an arbitrary function. Therefore, we have proved the following result.

Proposition 4.11. *For transversely isotropic linear elastic solids with the isotropy plane parallel to the x_1x_2 -plane in a Cartesian coordinate system (x_1, x_2, x_3) , the universal elastic strains have the following form*

$$\boldsymbol{\epsilon}^e(\mathbf{x}) = \begin{bmatrix} \epsilon_{11}^e(x_1, x_2, x_3) & \epsilon_{12}^e(x_1, x_2, x_3) & \epsilon_{13}^e(x_1, x_2) \\ \epsilon_{12}^e(x_1, x_2, x_3) & -\epsilon_{11}^e(x_1, x_2, x_3) + c_0 & \epsilon_{23}^e(x_1, x_2) \\ \epsilon_{13}^e(x_1, x_2) & \epsilon_{23}^e(x_1, x_2) & \epsilon_{33}^e \end{bmatrix},\tag{4.41}$$

³Note that there is a typo in [Yavari et al., 2020, Proposition 3.6].

where ϵ_{33}^e is constant. $\epsilon_{11}^e(x_1, x_2, x_3)$ satisfies the PDE (4.40)₁, and $\epsilon_{12}^e(x_1, x_2, x_3)$ has the representation (4.40)₂. $\epsilon_{13}^e(x_1, x_2)$ is an arbitrary function, while $\epsilon_{23}^e(x_1, x_2)$ has the representation (4.38). Universal eigenstrains (up to impotent eigenstrains) are the set of solutions of the six second-order linear PDEs $\text{curl} \circ \text{curl} \epsilon^* = -\mathbf{R}^e$, where the incompatibility tensor of the universal elastic strains reads

$$\mathbf{R}^e = \begin{bmatrix} -\frac{\partial^2 \epsilon_{11}^e}{\partial x_3^2} & \frac{\partial^2 \epsilon_{12}^e}{\partial x_3^2} & \frac{\partial}{\partial x_2} \left(\frac{\partial \epsilon_{23}^e}{\partial x_1} - \frac{\partial \epsilon_{13}^e}{\partial x_2} \right) \\ \frac{\partial^2 \epsilon_{12}^e}{\partial x_3^2} & \frac{\partial^2 \epsilon_{11}^e}{\partial x_3^2} & \frac{\partial}{\partial x_1} \left(\frac{\partial \epsilon_{13}^e}{\partial x_2} - \frac{\partial \epsilon_{23}^e}{\partial x_1} \right) \\ \frac{\partial}{\partial x_2} \left(\frac{\partial \epsilon_{23}^e}{\partial x_1} - \frac{\partial \epsilon_{13}^e}{\partial x_2} \right) & \frac{\partial}{\partial x_1} \left(\frac{\partial \epsilon_{13}^e}{\partial x_2} - \frac{\partial \epsilon_{23}^e}{\partial x_1} \right) & 0 \end{bmatrix}. \quad (4.42)$$

4.7 Cubic solids

A cubic solid has nine planes of symmetry at every point such that their normals are parallel to the edges and face diagonals of a cube. Suppose the edges of the cube are parallel to the coordinate axes of a Cartesian coordinate system (x_1, x_2, x_3) . A cubic solid has 3 independent elastic constants and with respect to this coordinate system has an elasticity matrix with the following representation

$$\mathbf{c} = \begin{bmatrix} c_{11} & c_{12} & c_{12} & 0 & 0 & 0 \\ c_{12} & c_{11} & c_{12} & 0 & 0 & 0 \\ c_{12} & c_{12} & c_{11} & 0 & 0 & 0 \\ 0 & 0 & 0 & c_{44} & 0 & 0 \\ 0 & 0 & 0 & 0 & c_{44} & 0 \\ 0 & 0 & 0 & 0 & 0 & c_{44} \end{bmatrix}. \quad (4.43)$$

In [Yavari et al., 2020] it was shown that for cubic solids and in the absence of eigenstrains universal displacements have the following form

$$\begin{aligned} u_1(x_1, x_2, x_3) &= \frac{a}{2} x_1 (x_3^2 - x_2^2) + c_1 x_1 x_3 + b_1 x_1 x_2 + d_1 x_1 + g_1(x_2, x_3), \\ u_2(x_1, x_2, x_3) &= \frac{a}{2} x_2 (x_1^2 - x_3^2) + a_1 x_1 x_2 - c_1 x_2 x_3 + d_2 x_2 + g_2(x_1, x_3), \\ u_3(x_1, x_2, x_3) &= \frac{a}{2} x_3 (x_2^2 - x_1^2) - a_1 x_1 x_3 - b_1 x_2 x_3 + d_3 x_3 + g_3(x_1, x_2), \end{aligned} \quad (4.44)$$

where g_1 , g_2 , and g_3 are arbitrary harmonic functions.

The universality constraints (4.2) give us the following nine PDEs:

$$\begin{aligned} \frac{\partial \epsilon_{11}^e}{\partial x_1} - \frac{\partial \epsilon_{22}^e}{\partial x_2} - \frac{\partial \epsilon_{33}^e}{\partial x_3} &= 0, \\ \frac{\partial \epsilon_{22}^e}{\partial x_1} + \frac{\partial \epsilon_{33}^e}{\partial x_1} &= 0, \\ \frac{\partial \epsilon_{11}^e}{\partial x_2} + \frac{\partial \epsilon_{33}^e}{\partial x_2} &= 0, \\ \frac{\partial \epsilon_{11}^e}{\partial x_3} + \frac{\partial \epsilon_{22}^e}{\partial x_3} &= 0, \end{aligned} \quad (4.45)$$

and

$$\begin{aligned}
\frac{\partial \epsilon_{12}^e}{\partial x_1} + \frac{\partial \epsilon_{23}^e}{\partial x_3} &= 0, \\
\frac{\partial \epsilon_{12}^e}{\partial x_2} + \frac{\partial \epsilon_{13}^e}{\partial x_3} &= 0, \\
\frac{\partial \epsilon_{13}^e}{\partial x_1} + \frac{\partial \epsilon_{23}^e}{\partial x_2} &= 0.
\end{aligned} \tag{4.46}$$

From (4.45)₁, $\epsilon_{11}^e = \epsilon_{11}^e(x_2, x_3)$, $\epsilon_{22}^e = \epsilon_{22}^e(x_1, x_3)$, and $\epsilon_{33}^e = \epsilon_{33}^e(x_1, x_2)$. From the remaining PDEs in (4.45) one concludes that

$$\begin{aligned}
\epsilon_{11}^e(x_2, x_3) &= g(x_2) + h(x_3), \\
\epsilon_{22}^e(x_1, x_3) &= f(x_1) - h(x_3), \\
\epsilon_{33}^e(x_1, x_2) &= -f(x_1) - g(x_2),
\end{aligned} \tag{4.47}$$

where $f(x_1)$, $g(x_2)$, and $h(x_3)$ are arbitrary functions.

From (4.46) one concludes that

$$\begin{aligned}
\epsilon_{12}^e(x_1, x_2, x_3) &= f_3(x_1, x_3) + g_3(x_2, x_3), \\
\epsilon_{23}^e(x_1, x_2, x_3) &= f_1(x_1, x_2) + g_1(x_1, x_3), \\
\epsilon_{13}^e(x_1, x_2, x_3) &= f_2(x_1, x_2) + g_2(x_2, x_3),
\end{aligned} \tag{4.48}$$

such that

$$\begin{aligned}
\frac{\partial}{\partial x_1} f_3(x_1, x_3) + \frac{\partial}{\partial x_3} g_1(x_1, x_3) &= 0, \\
\frac{\partial}{\partial x_2} g_3(x_2, x_3) + \frac{\partial}{\partial x_3} g_2(x_2, x_3) &= 0, \\
\frac{\partial}{\partial x_1} f_2(x_1, x_2) + \frac{\partial}{\partial x_2} f_1(x_1, x_2) &= 0.
\end{aligned} \tag{4.49}$$

Thus

$$\begin{aligned}
g_1(x_1, x_3) &= - \int f_{3,1}(x_1, x_3) dx_3 + \alpha(x_1), \\
g_3(x_2, x_3) &= - \int g_{2,3}(x_2, x_3) dx_2 + \gamma(x_3), \\
f_2(x_1, x_2) &= - \int f_{1,2}(x_1, x_2) dx_1 + \beta(x_2),
\end{aligned} \tag{4.50}$$

where $\alpha(x_1)$, $\beta(x_2)$, and $\gamma(x_3)$ are arbitrary functions. Therefore, we have proved the following result.

Proposition 4.12. *For cubic linear elastic solids the universal elastic strains have the following form*

$$\epsilon^e(\mathbf{x}) = \begin{bmatrix} g(x_2) + h(x_3) & f_3(x_1, x_3) + g_3(x_2, x_3) & f_2(x_1, x_2) + g_2(x_2, x_3) \\ f_3(x_1, x_3) + g_3(x_2, x_3) & f(x_1) - h(x_3) & f_1(x_1, x_2) + g_1(x_1, x_3) \\ f_2(x_1, x_2) + g_2(x_2, x_3) & f_1(x_1, x_2) + g_1(x_1, x_3) & g(x_2) + h(x_3) \end{bmatrix}, \tag{4.51}$$

where $g_1(x_1, x_3)$, $g_3(x_2, x_3)$, and $f_2(x_1, x_2)$ are given in (4.50), and the remaining functions are arbitrary. Universal eigenstrains (up to impotent eigenstrains) are the set of solutions of the six second-order linear

PDEs $\text{curl} \circ \text{curl} \boldsymbol{\epsilon}^* = -\mathbf{R}^e$, where the incompatibility tensor of the universal elastic strains reads

$$\mathbf{R}^e = \begin{bmatrix} -g''(x_2) - h''(x_3) & -\nabla^2 f_3(x_1, x_3) - \nabla^2 g_3(x_2, x_3) & -\nabla^2 f_1(x_1, x_2) - \nabla^2 g_1(x_1, x_3) \\ -\nabla^2 f_3(x_1, x_3) - \nabla^2 g_3(x_2, x_3) & -f''(x_1) + h''(x_3) & -\nabla^2 f_2(x_1, x_2) - \nabla^2 g_2(x_2, x_3) \\ -\nabla^2 f_1(x_1, x_2) - \nabla^2 g_1(x_1, x_3) & -\nabla^2 f_2(x_1, x_2) - \nabla^2 g_2(x_2, x_3) & g''(x_2) + h''(x_3) \end{bmatrix}. \quad (4.52)$$

5 Conclusion

We have studied the universality of elastic and anelastic strains in anisotropic linear anelasticity. Universal displacements are those displacement fields that satisfy the equilibrium equations in the absence of body forces for arbitrary elastic constants in a given symmetry class. The universality constraints of linear anelasticity restrict the possible forms of elastic strains. We completely characterized the universal elastic strains for all the eight symmetry classes. We observed that for triclinic, monoclinic, and trigonal solids universal elastic strains are compatible. The total strain $\boldsymbol{\epsilon} = \boldsymbol{\epsilon}^e + \boldsymbol{\epsilon}^*$ is compatible, and hence, $\text{curl} \circ \text{curl} \boldsymbol{\epsilon}^e + \text{curl} \circ \text{curl} \boldsymbol{\epsilon}^* = \mathbf{0}$. Having determined the set of universal elastic strains for every symmetry class, the corresponding universal eigenstrains are found to be those that satisfy the linear second-order PDEs $\text{curl} \circ \text{curl} \boldsymbol{\epsilon}^* = -\text{curl} \circ \text{curl} \boldsymbol{\epsilon}^e$ for at least one universal elastic strain field $\boldsymbol{\epsilon}^e(\mathbf{x})$. For triclinic, monoclinic, and trigonal classes we showed that only compatible eigenstrains are universal. If universal eigenstrains are defined modulo the compatible eigenstrains, for these three classes the universal displacements are identical to those of the corresponding linear elasticity universal displacements that were characterized in [Yavari et al., 2020]. For the other five classes universal eigenstrains are solutions to a system of inhomogeneous PDEs with forcing terms that are certain arbitrary functions depending on the symmetry class. We observed that the smaller the symmetry group, the smaller the space of universal elastic strains, and consequently, the smaller the space of universal eigenstrains. Hence, we have achieved a complete classification of universal elastic strains in linear anelasticity.

Acknowledgments

Arash Yavari was supported by ARO W911NF-18-1-0003 and NSF – Grant No. CMMI 1939901. This work was also supported by the Engineering and Physical Sciences Research Council grant EP/R020205/1 to Alain Goriely.

References

- D. Ambrosi, M. Ben Amar, C. J. Cyron, A. DeSimone, A. Goriely, J. D. Humphrey, and E. Kuhl. Growth and remodelling of living tissues: perspectives, challenges and opportunities. *Journal of the Royal Society Interface*, 16(157):20190233, 2019.
- E. Beltrami. Osservazioni sulla nota precedente. *Atti Accad. Lincei Rend*, 1(5):141–142, 1892.
- P. Chadwick, M. Vianello, and S. C. Cowin. A new proof that the number of linear elastic symmetries is eight. *Journal of the Mechanics and Physics of Solids*, 49(11):2471–2492, 2001.
- S. C. Cowin and S. B. Doty. *Tissue Mechanics*. Springer Science & Business Media, 2007.
- S. C. Cowin and M. M. Mehrabadi. Anisotropic symmetries of linear elasticity. *Applied Mechanics Reviews*, 48(5):247–285, 1995.

- J. L. Diani and D. Parks. Problem of an inclusion in an infinite body, approach in large deformation. *Mechanics of materials*, 32(1):43–55, 2000.
- C. Eckart. The thermodynamics of irreversible processes. IV. The theory of elasticity and anelasticity. *Physical Review*, 73(4):373, 1948.
- J. L. Ericksen. Deformations possible in every isotropic, incompressible, perfectly elastic body. *Zeitschrift für Angewandte Mathematik und Physik (ZAMP)*, 5(6):466–489, 1954.
- J. L. Ericksen. Deformations possible in every compressible, isotropic, perfectly elastic material. *Journal of Mathematics and Physics*, 34(1-4):126–128, 1955.
- J. L. Ericksen and R. S. Rivlin. Large elastic deformations of homogeneous anisotropic materials. *Journal of Rational Mechanics and Analysis*, 3:281–301, 1954.
- J. Eshelby. The determination of the elastic field of an ellipsoidal inclusion, and related problems. In *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, volume 241, pages 376–396. The Royal Society, 1957.
- R. L. Fosdick. Remarks on Compatibility. *Modern Developments in the Mechanics of Continua*, pages 109–127, 1966.
- A. Golgoon and A. Yavari. On the stress field of a nonlinear elastic solid torus with a toroidal inclusion. *Journal of Elasticity*, 128(1):115–145, 2017.
- A. Golgoon and A. Yavari. Nonlinear elastic inclusions in anisotropic solids. *Journal of Elasticity*, 130(2):239–269, 2018.
- A. Golgoon and A. Yavari. On Hashin’s hollow cylinder and sphere assemblages in anisotropic nonlinear elasticity. *Submitted*, 2021.
- A. Golgoon, S. Sadik, and A. Yavari. Circumferentially-symmetric finite eigenstrains in incompressible isotropic nonlinear elastic wedges. *International Journal of Non-Linear Mechanics*, 84:116–129, 2016.
- C. Goodbrake, A. Goriely, and A. Yavari. The mathematical foundations of anelasticity: Existence of smooth global intermediate configurations. *Proceedings of the Royal Society A*, 2020a.
- C. Goodbrake, A. Yavari, and A. Goriely. The anelastic Ericksen problem: Universal deformations and universal eigenstrains in incompressible nonlinear anelasticity. *Journal of Elasticity*, 142(2):291–381, 2020b.
- M. E. Gurtin. A generalization of the Beltrami stress functions in continuum mechanics. *Archive for Rational Mechanics and Analysis*.
- M. E. Gurtin. The linear theory of elasticity. In *Handbuch der Physik, Band VIa/2*. Springer-Verlag, Berlin, 1972.
- T.-S. Jun and A. M. Korsunsky. Evaluation of residual stresses and strains using the eigenstrain reconstruction method. *International Journal of Solids and Structures*, 47(13):1678–1686, 2010.
- C. Kim and P. Schiavone. A circular inhomogeneity subjected to non-uniform remote loading in finite plane elastostatics. *International Journal of Non-Linear Mechanics*, 42(8):989–999, 2007.
- C. Kim and P. Schiavone. Designing an inhomogeneity with uniform interior stress in finite plane elastostatics. *Acta Mechanica*, 197(3-4):285–299, 2008.
- C. Kim, M. Vasudevan, and P. Schiavone. Eshelby’s conjecture in finite plane elastostatics. *The Quarterly Journal of Mechanics and Applied Mathematics*, 61(1):63–73, 2008.
- N. Kinoshita and T. Mura. Elastic fields of inclusions in anisotropic media. *Physica Status Solidi (a)*, 5(3):759–768, 1971.

- W. W. Klingbeil and R. T. Shield. On a class of solutions in plane finite elasticity. *Zeitschrift für angewandte Mathematik und Physik ZAMP*, 17(4):489–511, 1966.
- K. Kondo. A proposal of a new theory concerning the yielding of materials based on riemannian geometry. *The Journal of the Japan Society of Aeronautical Engineering*, 2(8):29–31, 1949.
- S. Li, R. A. Sauer, and G. Wang. The eshelby tensors in a finite spherical domain—part i: Theoretical formulations. 2007.
- R. D. Mindlin and D. H. Cheng. Nuclei of strain in the semi-infinite solid. *Journal of Applied Physics*, 21(9):926–930, 1950.
- T. Mura. *Micromechanics of Defects in Solids*. Martinus Nijhoff, 1982.
- H. Reissner. Eigenspannungen und eigenspannungsquellen. *Zeitschrift für Angewandte Mathematik und Mechanik*, 11(1):1–8, 1931.
- R. S. Rivlin. Large elastic deformations of isotropic materials IV. Further developments of the general theory. *Philosophical Transactions of the Royal Society of London A*, 241(835):379–397, 1948.
- R. S. Rivlin. Large elastic deformations of isotropic materials. V. the problem of flexure. *Proceedings of the Royal Society of London A*, 195(1043):463–473, 1949a.
- R. S. Rivlin. A note on the torsion of an incompressible highly elastic cylinder. In *Mathematical Proceedings of the Cambridge Philosophical Society*, volume 45, pages 485–487. Cambridge University Press, 1949b.
- C. Ru, P. Schiavone, L. Sudak, and A. Mioduchowski. Uniformity of stresses inside an elliptic inclusion in finite plane elastostatics. *International Journal of Non-Linear Mechanics*, 40(2):281–287, 2005.
- C.-Q. Ru and P. Schiavone. On the elliptic inclusion in anti-plane shear. *Mathematics and Mechanics of Solids*, 1(3):327–333, 1996.
- S. Sadik and A. Yavari. On the origins of the idea of the multiplicative decomposition of the deformation gradient. *Mathematics and Mechanics of Solids*, 22:771–772, 2017.
- M. Singh and A. C. Pipkin. Note on Ericksen’s problem. *Zeitschrift für angewandte Mathematik und Physik ZAMP*, 16(5):706–709, 1965.
- F. Sozio and A. Yavari. Riemannian and Euclidean material structures in anelasticity. *Mathematics and Mechanics of Solids*, 25(6):1267–1293, 2020.
- T. C. T. Ting. Generalized Cowin–Mehrabadi theorems and a direct proof that the number of linear elastic symmetries is eight. *International Journal of Solids and Structures*, 40(25):7129–7142, 2003.
- C. Truesdell. *The Elements of Continuum Mechanics*. Springer-Verlag, 1966.
- Y. Ueda, K. Fukuda, K. Nakacho, and S. Endo. A new measuring method of residual stresses with the aid of finite element method and reliability of estimated values. *Transactions of JWRI*, 4(2):123–131, 1975.
- A. Yavari. Compatibility equations of nonlinear elasticity for non-simply-connected bodies. *Archive for Rational Mechanics and Analysis*, 209(1):237–253, 2013.
- A. Yavari. Universal deformations in inhomogeneous isotropic nonlinear elastic solids. *Proceedings of the Royal Society A*, 477(2253):20210547, 2021a.
- A. Yavari. On Eshelby’s inclusion problem in nonlinear anisotropic elasticity. *Journal of Micromechanics and Molecular Physics*, 6(1):2150002, 2021b.
- A. Yavari and A. Goriely. On the stress singularities generated by anisotropic eigenstrains and the hydrostatic stress due to annular inhomogeneities. *Journal of the Mechanics and Physics of Solids*, 76:325–337, 2015a.

- A. Yavari and A. Goriely. The twist-fit problem: Finite torsional and shear eigenstrains in nonlinear elastic solids. *Proceedings of the Royal Society of London A*, 471(2183), 2015b.
- A. Yavari and A. Goriely. The anelastic Ericksen problem: Universal eigenstrains and deformations in compressible isotropic elastic solids. *Proceedings of the Royal Society A*, 472(2196):20160690, 2016.
- A. Yavari and A. Goriely. Universal deformations in anisotropic nonlinear elastic solids. *Journal of the Mechanics and Physics of Solids*, 156:104598, 2021.
- A. Yavari and A. Goriely. The universal program of nonlinear hyperelasticity. *Journal of Elasticity*, 2022a.
- A. Yavari and A. Goriely. The universal program of linear elasticity. *Mathematics and Mechanics of Solids*, pages 1–18, 2022b. doi: 10.1177/10812865221091305.
- A. Yavari, C. Goodbrake, and A. Goriely. Universal displacements in linear elasticity. *Journal of the Mechanics and Physics of Solids*, 135:103782, 2020.
- K. Zhou, H. J. Hoh, X. Wang, L. M. Keer, J. H. Pang, B. Song, and Q. J. Wang. A review of recent works on inclusions. *Mechanics of Materials*, 60:144–158, 2013.